

表 532 専従専門医の有無 %クレアチニン産生速度(%)別(施設血液透析患者)

%クレアチニン 産生速度	なし	あり	合計	記載なし	総計
60未満	14,822	15,680	30,502		30,502
(%)	(48.6)	(51.4)	(100.0)		
60～	7,391	8,659	16,050		16,050
(%)	(46.0)	(54.0)	(100.0)		
70～	9,499	11,318	20,817		20,817
(%)	(45.6)	(54.4)	(100.0)		
80～	11,330	14,533	25,863		25,863
(%)	(43.8)	(56.2)	(100.0)		
90～	13,009	17,055	30,064		30,064
(%)	(43.3)	(56.7)	(100.0)		
100～	12,887	17,739	30,626		30,626
(%)	(42.1)	(57.9)	(100.0)		
110～	10,953	15,531	26,484		26,484
(%)	(41.4)	(58.6)	(100.0)		
120～	7,611	11,141	18,752		18,752
(%)	(40.6)	(59.4)	(100.0)		
130～	4,253	6,073	10,326		10,326
(%)	(41.2)	(58.8)	(100.0)		
140～	2,936	4,127	7,063		7,063
(%)	(41.6)	(58.4)	(100.0)		
合計	94,691	121,856	216,547		216,547
(%)	(43.7)	(56.3)	(100.0)		
記載なし	15,801	23,293	39,094		39,094
(%)	(40.4)	(59.6)	(100.0)		
総計	110,492	145,149	255,641		255,641
(%)	(43.2)	(56.8)	(100.0)		
平均	90.74	93.70	92.41		92.41
標準偏差	29.50	28.60	29.03		29.03

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査