

表 533 専従専門医の有無 体重減少量(kg)別(施設血液透析患者)

体重減少率	なし	あり	合計	記載なし	総計
1.0未満	3,911	4,462	8,373		8,373
(%)	(46.7)	(53.3)	(100.0)		
1.0～	5,012	6,507	11,519		11,519
(%)	(43.5)	(56.5)	(100.0)		
2.0～	10,784	14,650	25,434		25,434
(%)	(42.4)	(57.6)	(100.0)		
3.0～	19,981	27,396	47,377		47,377
(%)	(42.2)	(57.8)	(100.0)		
4.0～	25,517	34,015	59,532		59,532
(%)	(42.9)	(57.1)	(100.0)		
5.0～	19,366	25,337	44,703		44,703
(%)	(43.3)	(56.7)	(100.0)		
6.0～	9,808	12,341	22,149		22,149
(%)	(44.3)	(55.7)	(100.0)		
7.0～	3,780	4,689	8,469		8,469
(%)	(44.6)	(55.4)	(100.0)		
8.0～	1,250	1,376	2,626		2,626
(%)	(47.6)	(52.4)	(100.0)		
9.0～	676	813	1,489		1,489
(%)	(45.4)	(54.6)	(100.0)		
合計	100,085	131,586	231,671		231,671
(%)	(43.2)	(56.8)	(100.0)		
記載なし	10,407	13,563	23,970		23,970
(%)	(43.4)	(56.6)	(100.0)		
総計	110,492	145,149	255,641		255,641
(%)	(43.2)	(56.8)	(100.0)		
平均	4.36	4.34	4.35		4.35
標準偏差	1.76	1.71	1.73		1.73

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査