

表 534 専従専門医の有無 身長(cm)別(腹膜透析患者)

身長	なし	あり	合計	記載なし	総計
130未満	34	27	61	1	62
(%)	(55.7)	(44.3)	(100.0)		
130～	8	3	11		11
(%)	(72.7)	(27.3)	(100.0)		
135～	31	31	62		62
(%)	(50.0)	(50.0)	(100.0)		
140～	99	128	227		227
(%)	(43.6)	(56.4)	(100.0)		
145～	191	279	470		470
(%)	(40.6)	(59.4)	(100.0)		
150～	334	533	867		867
(%)	(38.5)	(61.5)	(100.0)		
155～	406	643	1,049		1,049
(%)	(38.7)	(61.3)	(100.0)		
160～	515	815	1,330		1,330
(%)	(38.7)	(61.3)	(100.0)		
165～	440	772	1,212		1,212
(%)	(36.3)	(63.7)	(100.0)		
170～	339	565	904		904
(%)	(37.5)	(62.5)	(100.0)		
175～	139	227	366		366
(%)	(38.0)	(62.0)	(100.0)		
180～	33	54	87		87
(%)	(37.9)	(62.1)	(100.0)		
185～	4	9	13		13
(%)	(30.8)	(69.2)	(100.0)		
190～	2	2	4		4
(%)	(50.0)	(50.0)	(100.0)		
合計	2,575	4,088	6,663	1	6,664
(%)	(38.6)	(61.4)	(100.0)		
記載なし	814	1,463	2,277		2,277
(%)	(35.7)	(64.3)	(100.0)		
総計	3,389	5,551	8,940	1	8,941
(%)	(37.9)	(62.1)	(100.0)		
平均	159.99	160.93	160.57	97.00	160.56
標準偏差	12.28	11.08	11.57		11.59

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査