

表 535 専従専門医の有無 体重(kg)別(腹膜透析患者)

体重	なし	あり	合計	記載なし	総計
20未満	17	21	38		38
(%)	(44.7)	(55.3)	(100.0)		
20～	15	7	22	1	23
(%)	(68.2)	(31.8)	(100.0)		
30～	77	95	172		172
(%)	(44.8)	(55.2)	(100.0)		
40～	359	545	904		904
(%)	(39.7)	(60.3)	(100.0)		
50～	645	946	1,591		1,591
(%)	(40.5)	(59.5)	(100.0)		
60～	614	1,002	1,616		1,616
(%)	(38.0)	(62.0)	(100.0)		
70～	320	491	811		811
(%)	(39.5)	(60.5)	(100.0)		
80～	108	190	298		298
(%)	(36.2)	(63.8)	(100.0)		
90～	36	68	104		104
(%)	(34.6)	(65.4)	(100.0)		
100～	10	10	20		20
(%)	(50.0)	(50.0)	(100.0)		
110～	5	5	10		10
(%)	(50.0)	(50.0)	(100.0)		
120～	2	7	9		9
(%)	(22.2)	(77.8)	(100.0)		
合計	2,208	3,387	5,595	1	5,596
(%)	(39.5)	(60.5)	(100.0)		
記載なし	1,181	2,164	3,345		3,345
(%)	(35.3)	(64.7)	(100.0)		
総計	3,389	5,551	8,940	1	8,941
(%)	(37.9)	(62.1)	(100.0)		
平均	60.26	61.07	60.75	21.30	60.74
標準偏差	14.09	13.74	13.89		13.90

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査