

表 536 専従専門医の有無 BUN(mg/dL)別(腹膜透析患者)

BUN	なし	あり	合計	記載なし	総計
20未満	13	16	29		29
(%)	(44.8)	(55.2)	(100.0)		
20～	91	92	183		183
(%)	(49.7)	(50.3)	(100.0)		
30～	282	390	672		672
(%)	(42.0)	(58.0)	(100.0)		
40～	545	846	1,391	1	1,392
(%)	(39.2)	(60.8)	(100.0)		
50～	654	959	1,613		1,613
(%)	(40.5)	(59.5)	(100.0)		
60～	441	709	1,150		1,150
(%)	(38.3)	(61.7)	(100.0)		
70～	211	338	549		549
(%)	(38.4)	(61.6)	(100.0)		
80～	109	136	245		245
(%)	(44.5)	(55.5)	(100.0)		
90～	37	52	89		89
(%)	(41.6)	(58.4)	(100.0)		
100～	17	23	40		40
(%)	(42.5)	(57.5)	(100.0)		
110～	1	6	7		7
(%)	(14.3)	(85.7)	(100.0)		
120～	4	3	7		7
(%)	(57.1)	(42.9)	(100.0)		
合計	2,405	3,570	5,975	1	5,976
(%)	(40.3)	(59.7)	(100.0)		
記載なし	984	1,981	2,965		2,965
(%)	(33.2)	(66.8)	(100.0)		
総計	3,389	5,551	8,940	1	8,941
(%)	(37.9)	(62.1)	(100.0)		
平均	54.74	55.02	54.91	45.00	54.90
標準偏差	16.64	15.25	15.82		15.82

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査