

表 539 専従専門医の有無 CRP濃度 (mg/dL) 別 (腹膜透析患者)

CRP濃度	なし	あり	合計	記載なし	総計
0.2未満	1,185	1,881	3,066	1	3,067
(%)	(38.6)	(61.4)	(100.0)		
0.2～	366	629	995		995
(%)	(36.8)	(63.2)	(100.0)		
0.5～	186	263	449		449
(%)	(41.4)	(58.6)	(100.0)		
1.0～	122	186	308		308
(%)	(39.6)	(60.4)	(100.0)		
2.0～	96	106	202		202
(%)	(47.5)	(52.5)	(100.0)		
4.0～	37	46	83		83
(%)	(44.6)	(55.4)	(100.0)		
6.0～	19	21	40		40
(%)	(47.5)	(52.5)	(100.0)		
8.0～	7	17	24		24
(%)	(29.2)	(70.8)	(100.0)		
10.0～	18	16	34		34
(%)	(52.9)	(47.1)	(100.0)		
15.0～	3	4	7		7
(%)	(42.9)	(57.1)	(100.0)		
合計	2,039	3,169	5,208	1	5,209
(%)	(39.2)	(60.8)	(100.0)		
記載なし	1,350	2,382	3,732		3,732
(%)	(36.2)	(63.8)	(100.0)		
総計	3,389	5,551	8,940	1	8,941
(%)	(37.9)	(62.1)	(100.0)		
平均	0.70	0.58	0.63	0.03	0.63
標準偏差	1.93	1.55	1.71		1.71

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査