

表 540 専従専門医の有無 補正カルシウム濃度 (mg/dL) 別 (腹膜透析患者)

補正 カルシウム濃度	なし	あり	合計	記載なし	総計
7.0未満	13	16	29		29
(%)	(44.8)	(55.2)	(100.0)		
7.0～	20	16	36		36
(%)	(55.6)	(44.4)	(100.0)		
7.5～	44	83	127		127
(%)	(34.6)	(65.4)	(100.0)		
8.0～	116	170	286		286
(%)	(40.6)	(59.4)	(100.0)		
8.4～	34	52	86		86
(%)	(39.5)	(60.5)	(100.0)		
8.5～	368	538	906		906
(%)	(40.6)	(59.4)	(100.0)		
9.0～	611	991	1,602		1,602
(%)	(38.1)	(61.9)	(100.0)		
9.5～	646	983	1,629		1,629
(%)	(39.7)	(60.3)	(100.0)		
10.0～	88	103	191		191
(%)	(46.1)	(53.9)	(100.0)		
10.1～	221	335	556	1	557
(%)	(39.7)	(60.3)	(100.0)		
10.5～	124	172	296		296
(%)	(41.9)	(58.1)	(100.0)		
11.0～	49	62	111		111
(%)	(44.1)	(55.9)	(100.0)		
11.5～	17	12	29		29
(%)	(58.6)	(41.4)	(100.0)		
12.0～	15	12	27		27
(%)	(55.6)	(44.4)	(100.0)		
合計	2,366	3,545	5,911	1	5,912
(%)	(40.0)	(60.0)	(100.0)		
記載なし	1,023	2,006	3,029		3,029
(%)	(33.8)	(66.2)	(100.0)		
総計	3,389	5,551	8,940	1	8,941
(%)	(37.9)	(62.1)	(100.0)		
平均	9.42	9.40	9.41	10.30	9.41
標準偏差	0.83	0.78	0.80		0.80

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査