

表 541 専従専門医の有無 リン濃度(mg/dL)別(腹膜透析患者)

リン濃度	なし	あり	合計	記載なし	総計
3未満	64	62	126		126
(%)	(50.8)	(49.2)	(100.0)		
3.0～	117	153	270		270
(%)	(43.3)	(56.7)	(100.0)		
3.5～	228	297	525		525
(%)	(43.4)	(56.6)	(100.0)		
4.0～	296	473	769	1	770
(%)	(38.5)	(61.5)	(100.0)		
4.5～	334	587	921		921
(%)	(36.3)	(63.7)	(100.0)		
5.0～	389	559	948		948
(%)	(41.0)	(59.0)	(100.0)		
5.5～	335	504	839		839
(%)	(39.9)	(60.1)	(100.0)		
6.0～	42	96	138		138
(%)	(30.4)	(69.6)	(100.0)		
6.1～	179	298	477		477
(%)	(37.5)	(62.5)	(100.0)		
6.5～	167	232	399		399
(%)	(41.9)	(58.1)	(100.0)		
7.0～	91	121	212		212
(%)	(42.9)	(57.1)	(100.0)		
7.5～	64	85	149		149
(%)	(43.0)	(57.0)	(100.0)		
8.0～	35	36	71		71
(%)	(49.3)	(50.7)	(100.0)		
8.5～	13	25	38		38
(%)	(34.2)	(65.8)	(100.0)		
9.0～	44	45	89		89
(%)	(49.4)	(50.6)	(100.0)		
合計	2,398	3,573	5,971	1	5,972
(%)	(40.2)	(59.8)	(100.0)		
記載なし	991	1,978	2,969		2,969
(%)	(33.4)	(66.6)	(100.0)		
総計	3,389	5,551	8,940	1	8,941
(%)	(37.9)	(62.1)	(100.0)		
平均	5.25	5.25	5.25	4.00	5.25
標準偏差	1.42	1.30	1.35		1.35

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査