

表 543 専従専門医の有無 PTH値 (pg/mL) 別 (腹膜透析患者、PTH測定法:whole-PTH)

PTH値	なし	あり	合計	記載なし	総計
20未満	21	19	40		40
(%)	(52.5)	(47.5)	(100.0)		
20～	16	29	45		45
(%)	(35.6)	(64.4)	(100.0)		
35～	18	37	55		55
(%)	(32.7)	(67.3)	(100.0)		
50～	17	60	77		77
(%)	(22.1)	(77.9)	(100.0)		
75～	21	48	69		69
(%)	(30.4)	(69.6)	(100.0)		
100～	12	59	71		71
(%)	(16.9)	(83.1)	(100.0)		
125～	9	36	45		45
(%)	(20.0)	(80.0)	(100.0)		
150～		1	1		1
(%)		(100.0)	(100.0)		
151～	8	35	43		43
(%)	(18.6)	(81.4)	(100.0)		
180～	11	23	34		34
(%)	(32.4)	(67.6)	(100.0)		
210～	5	22	27		27
(%)	(18.5)	(81.5)	(100.0)		
240～	11	12	23		23
(%)	(47.8)	(52.2)	(100.0)		
300～	7	12	19		19
(%)	(36.8)	(63.2)	(100.0)		
350～	1	7	8		8
(%)	(12.5)	(87.5)	(100.0)		
400～	6	5	11		11
(%)	(54.5)	(45.5)	(100.0)		
500～	3	8	11		11
(%)	(27.3)	(72.7)	(100.0)		
合計	166	413	579		579
(%)	(28.7)	(71.3)	(100.0)		
記載なし	28	112	140		140
(%)	(20.0)	(80.0)	(100.0)		
総計	194	525	719		719
(%)	(27.0)	(73.0)	(100.0)		
平均	136.13	131.12	132.56		132.56
標準偏差	159.13	115.50	129.39		129.39

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査