

表 544 専従専門医の有無 ヘモグロビン濃度 (g/dL) 別 (腹膜透析患者)

ヘモグロビン 濃度	なし	あり	合計	記載なし	総計
5未満	3	4	7		7
(%)	(42.9)	(57.1)	(100.0)		
5～	5	9	14		14
(%)	(35.7)	(64.3)	(100.0)		
6～	19	9	28		28
(%)	(67.9)	(32.1)	(100.0)		
7～	62	61	123	1	124
(%)	(50.4)	(49.6)	(100.0)		
8～	164	180	344		344
(%)	(47.7)	(52.3)	(100.0)		
9～	386	517	903		903
(%)	(42.7)	(57.3)	(100.0)		
10～	685	1,021	1,706		1,706
(%)	(40.2)	(59.8)	(100.0)		
11～	651	1,058	1,709		1,709
(%)	(38.1)	(61.9)	(100.0)		
12～	286	510	796		796
(%)	(35.9)	(64.1)	(100.0)		
13～	98	139	237		237
(%)	(41.4)	(58.6)	(100.0)		
14～	36	43	79		79
(%)	(45.6)	(54.4)	(100.0)		
合計	2,395	3,551	5,946	1	5,947
(%)	(40.3)	(59.7)	(100.0)		
記載なし	994	2,000	2,994		2,994
(%)	(33.2)	(66.8)	(100.0)		
総計	3,389	5,551	8,940	1	8,941
(%)	(37.9)	(62.1)	(100.0)		
平均	10.73	10.87	10.81	7.30	10.81
標準偏差	1.47	1.38	1.42		1.42

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査