

表 2126 HDF希釈方法 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
施設血液透析							255,641	255,641
(%)							(94.0)	(82.4)
Off-line HDF	638	4,275	5	5	4,923		1,392	6,315
(%)	(2.1)	(72.4)	(100.0)	(0.3)	(12.9)		(0.5)	(2.0)
On-line HDF	29,881	1,342		1,317	32,540	1	3,549	36,090
(%)	(97.6)	(22.7)		(82.8)	(85.4)	(100.0)	(1.3)	(11.6)
Push/Pull HDF	8	88		266	362		175	537
(%)	(0.0)	(1.5)		(16.7)	(0.9)		(0.1)	(0.2)
ハイオフィльтраーション	2	154		1	157		184	341
(%)	(0.0)	(2.6)		(0.1)	(0.4)		(0.1)	(0.1)
血液濾過							92	92
(%)							(0.0)	(0.0)
血液吸着透析 (リクセル)	2	4		1	7		1,616	1,623
(%)	(0.0)	(0.1)		(0.1)	(0.0)		(0.6)	(0.5)
血液吸着透析 (その他)							7	7
(%)							(0.0)	(0.0)
在宅血液透析							521	521
(%)							(0.2)	(0.2)
腹膜透析 (CAPD)	41	15			56		4,965	5,021
(%)	(0.1)	(0.3)			(0.1)		(1.8)	(1.6)
腹膜透析 (APD)	25	15			40		2,799	2,839
(%)	(0.1)	(0.3)			(0.1)		(1.0)	(0.9)
腹膜透析 (CCPD)	15	13			28		1,053	1,081
(%)	(0.0)	(0.2)			(0.1)		(0.4)	(0.3)
合計	30,612	5,906	5	1,590	38,113	1	271,994	310,108
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
記載なし								
総計	30,612	5,906	5	1,590	38,113	1	271,994	310,108

数値下のかっこ内は列方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2014年調査