

表 3435 透析前補正カルシウム濃度(mg/dL) 透析前リン濃度(mg/dL)別 (透析患者全体 インタクトPTH値160～180)

透析前リン濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
3未満	3	2	6	23	6	93	115	55	5	20	11	5	4	2	350	5	355	9.18	0.85
(%)	(0.9)	(0.6)	(1.7)	(6.6)	(1.7)	(26.6)	(32.9)	(15.7)	(1.4)	(5.7)	(3.1)	(1.4)	(1.1)	(0.6)	(100.0)				
3.0～	1	2	12	58	11	208	196	94	8	27	19	2		1	639	1	640	9.08	0.64
(%)	(0.2)	(0.3)	(1.9)	(9.1)	(1.7)	(32.6)	(30.7)	(14.7)	(1.3)	(4.2)	(3.0)	(0.3)		(0.2)	(100.0)				
3.5～	3	8	22	120	32	372	394	229	15	40	18	9	1		1,263	13	1,276	9.06	0.63
(%)	(0.2)	(0.6)	(1.7)	(9.5)	(2.5)	(29.5)	(31.2)	(18.1)	(1.2)	(3.2)	(1.4)	(0.7)	(0.1)		(100.0)				
4.0～	5	9	36	144	58	589	635	327	42	92	51	10	1	2	2,001	21	2,022	9.11	0.66
(%)	(0.2)	(0.4)	(1.8)	(7.2)	(2.9)	(29.4)	(31.7)	(16.3)	(2.1)	(4.6)	(2.5)	(0.5)	(0.0)	(0.1)	(100.0)				
4.5～	3	14	51	175	62	702	779	452	52	113	57	12	8	2	2,482	26	2,508	9.13	0.66
(%)	(0.1)	(0.6)	(2.1)	(7.1)	(2.5)	(28.3)	(31.4)	(18.2)	(2.1)	(4.6)	(2.3)	(0.5)	(0.3)	(0.1)	(100.0)				
5.0～	5	6	58	207	60	765	781	430	55	128	48	12	8	1	2,564	22	2,586	9.10	0.65
(%)	(0.2)	(0.2)	(2.3)	(8.1)	(2.3)	(29.8)	(30.5)	(16.8)	(2.1)	(5.0)	(1.9)	(0.5)	(0.3)	(0.0)	(100.0)				
5.5～	12	10	48	163	53	638	648	388	31	128	51	7	5	1	2,183	16	2,199	9.11	0.70
(%)	(0.5)	(0.5)	(2.2)	(7.5)	(2.4)	(29.2)	(29.7)	(17.8)	(1.4)	(5.9)	(2.3)	(0.3)	(0.2)	(0.0)	(100.0)				
6.0～			8	39	15	99	124	51	8	15	9	2			370		370	9.08	0.62
(%)			(2.2)	(10.5)	(4.1)	(26.8)	(33.5)	(13.8)	(2.2)	(4.1)	(2.4)	(0.5)			(100.0)				
6.1～	2	8	29	112	29	321	374	237	16	65	28	2		2	1,225	12	1,237	9.11	0.66
(%)	(0.2)	(0.7)	(2.4)	(9.1)	(2.4)	(26.2)	(30.5)	(19.3)	(1.3)	(5.3)	(2.3)	(0.2)		(0.2)	(100.0)				
6.5～	6	8	32	122	28	304	273	182	23	49	22	13	3	1	1,066	8	1,074	9.06	0.76
(%)	(0.6)	(0.8)	(3.0)	(11.4)	(2.6)	(28.5)	(25.6)	(17.1)	(2.2)	(4.6)	(2.1)	(1.2)	(0.3)	(0.1)	(100.0)				
7.0～	3	6	15	66	18	166	161	97	9	40	18	3	1	1	604	7	611	9.07	0.75
(%)	(0.5)	(1.0)	(2.5)	(10.9)	(3.0)	(27.5)	(26.7)	(16.1)	(1.5)	(6.6)	(3.0)	(0.5)	(0.2)	(0.2)	(100.0)				
7.5～		6	16	39	18	128	102	61	8	14	5	3	2		402	7	409	8.99	0.69
(%)		(1.5)	(4.0)	(9.7)	(4.5)	(31.8)	(25.4)	(15.2)	(2.0)	(3.5)	(1.2)	(0.7)	(0.5)		(100.0)				
8.0～	4	3	8	30	2	57	48	43	1	14	4		2		216	3	219	9.00	0.84
(%)	(1.9)	(1.4)	(3.7)	(13.9)	(0.9)	(26.4)	(22.2)	(19.9)	(0.5)	(6.5)	(1.9)		(0.9)		(100.0)				
8.5～	1	2	3	14	5	37	44	21	2	5	2	1			137	2	139	9.02	0.72
(%)	(0.7)	(1.5)	(2.2)	(10.2)	(3.6)	(27.0)	(32.1)	(15.3)	(1.5)	(3.6)	(1.5)	(0.7)			(100.0)				
9.0～	3	2	15	25	8	46	53	45	5	11	5	2	1		221	2	223	9.02	0.83
(%)	(1.4)	(0.9)	(6.8)	(11.3)	(3.6)	(20.8)	(24.0)	(20.4)	(2.3)	(5.0)	(2.3)	(0.9)	(0.5)		(100.0)				
合計	51	86	359	1,337	405	4,525	4,727	2,712	280	761	348	83	36	13	15,723	145	15,868	9.09	0.68
(%)	(0.3)	(0.5)	(2.3)	(8.5)	(2.6)	(28.8)	(30.1)	(17.2)	(1.8)	(4.8)	(2.2)	(0.5)	(0.2)	(0.1)	(100.0)				
記載なし				1			2								3	22	25	8.80	0.44
(%)				(33.3)			(66.7)								(100.0)				
総計	51	86	359	1,338	405	4,525	4,729	2,712	280	761	348	83	36	13	15,726	167	15,893	9.09	0.68
(%)	(0.3)	(0.5)	(2.3)	(8.5)	(2.6)	(28.8)	(30.1)	(17.2)	(1.8)	(4.8)	(2.2)	(0.5)	(0.2)	(0.1)	(100.0)				
平均	5.93	5.66	5.64	5.42	5.40	5.25	5.20	5.31	5.29	5.37	5.22	5.28	5.43	4.75	5.28	5.34	5.28		
標準偏差	2.20	1.56	1.60	1.41	1.37	1.31	1.30	1.33	1.30	1.34	1.35	1.72	1.54	1.73	1.34	1.43	1.34		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査