

表 1951 透析時間(時間/回) 治療方法(新分類)別 (体外循環を用いた透析療法*)

治療方法 (新分類)	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
施設血液透析	699	27,588	18,566	158,787	11,862	15,866	624	1,049	61	557	235,659	19,982	255,641	3.96	0.53
(%)	(83.7)	(91.6)	(87.7)	(85.0)	(74.0)	(78.1)	(72.0)	(74.9)	(71.8)	(86.1)	(84.7)	(87.3)	(84.9)		
Off-line HDF	13	388	296	3,978	524	629	27	39	6	6	5,906	409	6,315	4.09	0.52
(%)	(1.6)	(1.3)	(1.4)	(2.1)	(3.3)	(3.1)	(3.1)	(2.8)	(7.1)	(0.9)	(2.1)	(1.8)	(2.1)		
On-line HDF	88	2,002	2,191	22,259	3,329	3,437	201	249	11	31	33,798	2,292	36,090	4.10	0.51
(%)	(10.5)	(6.6)	(10.4)	(11.9)	(20.8)	(16.9)	(23.2)	(17.8)	(12.9)	(4.8)	(12.1)	(10.0)	(12.0)		
Push/Pull HDF		48	42	384	15	21		3			513	24	537	3.94	0.42
(%)		(0.2)	(0.2)	(0.2)	(0.1)	(0.1)		(0.2)			(0.2)	(0.1)	(0.2)		
バイオフィльтраーション		12	16	220	42	32	1	2			325	16	341	4.13	0.45
(%)		(0.0)	(0.1)	(0.1)	(0.3)	(0.2)	(0.1)	(0.1)			(0.1)	(0.1)	(0.1)		
血液濾過	1	6		62	4	17	1	1			92		92	4.18	0.60
(%)	(0.1)	(0.0)		(0.0)	(0.0)	(0.1)	(0.1)	(0.1)			(0.0)		(0.0)		
血液吸着透析 (リクセル)		13	38	1,000	231	235	11	25	1	8	1,562	61	1,623	4.27	0.53
(%)		(0.0)	(0.2)	(0.5)	(1.4)	(1.2)	(1.3)	(1.8)	(1.2)	(1.2)	(0.6)	(0.3)	(0.5)		
血液吸着透析 (その他)				4							4	3	7	4.00	
(%)				(0.0)							(0.0)	(0.0)	(0.0)		
在宅血液透析	34	73	19	98	14	86	2	32	6	45	409	112	521	4.48	1.54
(%)	(4.1)	(0.2)	(0.1)	(0.1)	(0.1)	(0.4)	(0.2)	(2.3)	(7.1)	(7.0)	(0.1)	(0.5)	(0.2)		
合計	835	30,130	21,168	186,792	16,021	20,323	867	1,400	85	647	278,268	22,899	301,167	3.98	0.53
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		
記載なし															
総計	835	30,130	21,168	186,792	16,021	20,323	867	1,400	85	647	278,268	22,899	301,167	3.98	0.53

数値下のかっこ内は列方向の合計に対する%です。

*体外循環を用いた透析療法=施設血液透析、血液透析濾過、血液濾過、血液吸着透析、在宅血液透析

(C)Japanese Society for Dialysis Therapy

2014年調査