

表 1952 透析時間(時間/回) 治療方法別 (体外循環を用いた透析療法*)

治療方法	3.0未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.5～	7.0～	合計	記載なし	総計	平均	標準偏差
施設血液透析	699	27,588	18,566	158,787	11,862	15,866	624	1,049	61	557	235,659	19,982	255,641	3.96	0.53
(%)	(83.7)	(91.6)	(87.7)	(85.0)	(74.0)	(78.1)	(72.0)	(74.9)	(71.8)	(86.1)	(84.7)	(87.3)	(84.9)		
血液透析濾過	101	2,450	2,545	26,841	3,910	4,119	229	293	17	37	40,542	2,741	43,283	4.10	0.51
(%)	(12.1)	(8.1)	(12.0)	(14.4)	(24.4)	(20.3)	(26.4)	(20.9)	(20.0)	(5.7)	(14.6)	(12.0)	(14.4)		
血液濾過	1	6	62	4	17	1	1	1			92		92	4.18	0.60
(%)	(0.1)	(0.0)		(0.0)	(0.0)	(0.1)	(0.1)	(0.1)			(0.0)		(0.0)		
血液吸着透析		13	38	1,004	231	235	11	25	1	8	1,566	64	1,630	4.27	0.53
(%)		(0.0)	(0.2)	(0.5)	(1.4)	(1.2)	(1.3)	(1.8)	(1.2)	(1.2)	(0.6)	(0.3)	(0.5)		
在宅血液透析	34	73	19	98	14	86	2	32	6	45	409	112	521	4.48	1.54
(%)	(4.1)	(0.2)	(0.1)	(0.1)	(0.1)	(0.4)	(0.2)	(2.3)	(7.1)	(7.0)	(0.1)	(0.5)	(0.2)		
合計	835	30,130	21,168	186,792	16,021	20,323	867	1,400	85	647	278,268	22,899	301,167	3.98	0.53
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		
記載なし															
総計	835	30,130	21,168	186,792	16,021	20,323	867	1,400	85	647	278,268	22,899	301,167	3.98	0.53

数値下のかっこ内は列方向の合計に対する%です。
*体外循環を用いた透析療法=施設血液透析、血液透析濾過、血液濾過、血液吸着透析、在宅血液透析
(C)Japanese Society for Dialysis Therapy
2014年調査