

表 2752 体重減少率(%) 治療方法(新分類)別(体外循環を用いた透析療法* 透析回数週3回 透析歴2年以上)

治療方法 (新分類)	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
施設血液透析	3,696	6,609	16,921	34,584	46,415	36,289	18,427	7,108	2,195	1,205	173,449	4,760	178,209	4.53	1.65
(%)	(85.8)	(85.2)	(84.1)	(82.6)	(82.5)	(82.4)	(82.6)	(82.9)	(82.9)	(86.0)	(82.9)	(85.0)	(82.9)		
Off-line HDF	95	175	461	1,040	1,393	1,121	600	209	81	33	5,208	112	5,320	4.61	1.61
(%)	(2.2)	(2.3)	(2.3)	(2.5)	(2.5)	(2.5)	(2.7)	(2.4)	(3.1)	(2.4)	(2.5)	(2.0)	(2.5)		
On-line HDF	485	887	2,516	5,697	7,820	6,140	3,061	1,166	348	154	28,274	682	28,956	4.59	1.58
(%)	(11.3)	(11.4)	(12.5)	(13.6)	(13.9)	(13.9)	(13.7)	(13.6)	(13.1)	(11.0)	(13.5)	(12.2)	(13.5)		
Push/Pull HDF	5	19	33	99	110	87	29	9	3	1	395	12	407	4.36	1.44
(%)	(0.1)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.2)		
バイオフィルトレーション	2	10	20	57	54	76	35	24	4	2	284	4	288	4.87	1.69
(%)	(0.0)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.3)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)		
血液濾過	1	2	6	13	25	10	5	1		1	64		64	4.40	1.49
(%)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)		(0.1)	(0.0)		(0.0)		
血液吸着透析 (リクセル)	19	47	158	338	422	321	146	61	16	5	1,533	19	1,552	4.49	1.50
(%)	(0.4)	(0.6)	(0.8)	(0.8)	(0.8)	(0.7)	(0.7)	(0.7)	(0.6)	(0.4)	(0.7)	(0.3)	(0.7)		
血液吸着透析 (その他)				1	2						3	1	4	4.13	0.74
(%)				(0.0)	(0.0)						(0.0)	(0.0)	(0.0)		
在宅血液透析	3	5	16	16	12	8	3				63	8	71	3.49	1.49
(%)	(0.1)	(0.1)	(0.1)	(0.0)	(0.0)	(0.0)	(0.0)				(0.0)	(0.1)	(0.0)		
合計	4,306	7,754	20,131	41,845	56,253	44,052	22,306	8,578	2,647	1,401	209,273	5,598	214,871	4.54	1.64
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		
記載なし															
総計	4,306	7,754	20,131	41,845	56,253	44,052	22,306	8,578	2,647	1,401	209,273	5,598	214,871	4.54	1.64

数値下のかっこ内は列方向の合計に対する%です。

*体外循環を用いた透析療法=施設血液透析、血液透析濾過、血液濾過、血液吸着透析、在宅血液透析

(C)Japanese Society for Dialysis Therapy

2014年調査