

表 570 糖尿病の既往 身長(cm)別（透析患者全体）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	183	146	329		25	354
(%)	(55.6)	(44.4)	(100.0)			
130～	402	220	622	1	73	696
(%)	(64.6)	(35.4)	(100.0)			
135～	1,551	983	2,534	1	302	2,837
(%)	(61.2)	(38.8)	(100.0)			
140～	5,343	3,906	9,249	18	1,030	10,297
(%)	(57.8)	(42.2)	(100.0)			
145～	11,999	9,128	21,127	34	2,512	23,673
(%)	(56.8)	(43.2)	(100.0)			
150～	18,941	15,168	34,109	43	4,083	38,235
(%)	(55.5)	(44.5)	(100.0)			
155～	21,301	18,739	40,040	43	4,389	44,472
(%)	(53.2)	(46.8)	(100.0)			
160～	23,307	25,349	48,656	43	4,951	53,650
(%)	(47.9)	(52.1)	(100.0)			
165～	20,239	23,503	43,742	33	4,213	47,988
(%)	(46.3)	(53.7)	(100.0)			
170～	12,983	15,299	28,282	26	2,766	31,074
(%)	(45.9)	(54.1)	(100.0)			
175～	4,626	5,696	10,322	8	989	11,319
(%)	(44.8)	(55.2)	(100.0)			
180～	1,174	1,560	2,734	3	242	2,979
(%)	(42.9)	(57.1)	(100.0)			
185～	177	245	422	1	42	465
(%)	(41.9)	(58.1)	(100.0)			
190～	17	55	72		5	77
(%)	(23.6)	(76.4)	(100.0)			
合計	122,243	119,997	242,240	254	25,622	268,116
(%)	(50.5)	(49.5)	(100.0)			
記載なし	7,661	16,429	24,090	104	17,798	41,992
(%)	(31.8)	(68.2)	(100.0)			
総計	129,904	136,426	266,330	358	43,420	310,108
(%)	(48.8)	(51.2)	(100.0)			
平均	159.28	160.97	160.12	158.18	159.39	160.05
標準偏差	9.74	9.48	9.65	9.99	9.64	9.65

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査