

表 578 糖尿病の既往 PTH値(pg/mL)別（透析患者全体）PTH測定法:whole-PTH

| PTH値 | 既往なし   | 既往あり   | 合計      | 不明    | 記載なし   | 総計     |
|------|--------|--------|---------|-------|--------|--------|
| 20未満 | 833    | 898    | 1,731   |       | 163    | 1,894  |
| (%)  | (48.1) | (51.9) | (100.0) |       |        |        |
| 20～  | 1,054  | 1,091  | 2,145   | 1     | 185    | 2,331  |
| (%)  | (49.1) | (50.9) | (100.0) |       |        |        |
| 35～  | 945    | 917    | 1,862   | 1     | 180    | 2,043  |
| (%)  | (50.8) | (49.2) | (100.0) |       |        |        |
| 50～  | 1,437  | 1,437  | 2,874   |       | 279    | 3,153  |
| (%)  | (50.0) | (50.0) | (100.0) |       |        |        |
| 75～  | 1,269  | 1,215  | 2,484   |       | 264    | 2,748  |
| (%)  | (51.1) | (48.9) | (100.0) |       |        |        |
| 100～ | 1,029  | 1,047  | 2,076   |       | 222    | 2,298  |
| (%)  | (49.6) | (50.4) | (100.0) |       |        |        |
| 125～ | 703    | 647    | 1,350   |       | 189    | 1,539  |
| (%)  | (52.1) | (47.9) | (100.0) |       |        |        |
| 150～ | 21     | 22     | 43      |       | 5      | 48     |
| (%)  | (48.8) | (51.2) | (100.0) |       |        |        |
| 151～ | 604    | 518    | 1,122   | 1     | 140    | 1,263  |
| (%)  | (53.8) | (46.2) | (100.0) |       |        |        |
| 180～ | 429    | 372    | 801     |       | 74     | 875    |
| (%)  | (53.6) | (46.4) | (100.0) |       |        |        |
| 210～ | 292    | 278    | 570     |       | 58     | 628    |
| (%)  | (51.2) | (48.8) | (100.0) |       |        |        |
| 240～ | 349    | 256    | 605     |       | 69     | 674    |
| (%)  | (57.7) | (42.3) | (100.0) |       |        |        |
| 300～ | 145    | 101    | 246     |       | 38     | 284    |
| (%)  | (58.9) | (41.1) | (100.0) |       |        |        |
| 350～ | 92     | 71     | 163     |       | 16     | 179    |
| (%)  | (56.4) | (43.6) | (100.0) |       |        |        |
| 400～ | 107    | 69     | 176     |       | 26     | 202    |
| (%)  | (60.8) | (39.2) | (100.0) |       |        |        |
| 500～ | 142    | 75     | 217     |       | 30     | 247    |
| (%)  | (65.4) | (34.6) | (100.0) |       |        |        |
| 合計   | 9,451  | 9,014  | 18,465  | 3     | 1,938  | 20,406 |
| (%)  | (51.2) | (48.8) | (100.0) |       |        |        |
| 記載なし | 191    | 266    | 457     | 1     | 199    | 657    |
| (%)  | (41.8) | (58.2) | (100.0) |       |        |        |
| 総計   | 9,642  | 9,280  | 18,922  | 4     | 2,137  | 21,063 |
| (%)  | (51.0) | (49.0) | (100.0) |       |        |        |
| 平均   | 112.71 | 101.09 | 107.04  | 76.67 | 117.29 | 108.01 |
| 標準偏差 | 122.88 | 100.99 | 112.87  | 78.81 | 124.07 | 114.01 |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査