

表 579 糖尿病の既往 透析前ヘモグロビン濃度(g/dL)別（透析患者全体）

透析前ヘモグロビン濃度	既往なし	既往あり	合計	不明	記載なし	総計
5未満	168	121	289		14	303
(%)	(58.1)	(41.9)	(100.0)			
5～	167	234	401		28	429
(%)	(41.6)	(58.4)	(100.0)			
6～	472	575	1,047	5	108	1,160
(%)	(45.1)	(54.9)	(100.0)			
7～	1,923	2,084	4,007	7	421	4,435
(%)	(48.0)	(52.0)	(100.0)			
8～	6,365	6,819	13,184	22	1,348	14,554
(%)	(48.3)	(51.7)	(100.0)			
9～	20,472	21,049	41,521	54	4,414	45,989
(%)	(49.3)	(50.7)	(100.0)			
10～	42,975	41,750	84,725	90	9,097	93,912
(%)	(50.7)	(49.3)	(100.0)			
11～	35,979	34,503	70,482	69	7,622	78,173
(%)	(51.0)	(49.0)	(100.0)			
12～	13,286	12,908	26,194	24	2,636	28,854
(%)	(50.7)	(49.3)	(100.0)			
13～	3,245	3,379	6,624	4	680	7,308
(%)	(49.0)	(51.0)	(100.0)			
14～	1,187	1,254	2,441		232	2,673
(%)	(48.6)	(51.4)	(100.0)			
合計	126,239	124,676	250,915	275	26,600	277,790
(%)	(50.3)	(49.7)	(100.0)			
記載なし	3,665	11,750	15,415	83	16,820	32,318
(%)	(23.8)	(76.2)	(100.0)			
総計	129,904	136,426	266,330	358	43,420	310,108
(%)	(48.8)	(51.2)	(100.0)			
平均	10.72	10.70	10.71	10.41	10.71	10.71
標準偏差	1.27	1.30	1.29	1.31	1.25	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査