

表 581 糖尿病の既往 HDL-C濃度(mg/dL)別（透析患者全体）

HDL-C濃度	既往なし	既往あり	合計	不明	記載なし	総計
20未満	257	483	740		56	796
(%)	(34.7)	(65.3)	(100.0)			
20～	4,734	9,825	14,559	12	1,052	15,623
(%)	(32.5)	(67.5)	(100.0)			
30～	17,554	24,611	42,165	40	3,642	45,847
(%)	(41.6)	(58.4)	(100.0)			
40～	23,372	23,261	46,633	51	4,668	51,352
(%)	(50.1)	(49.9)	(100.0)			
50～	18,934	14,623	33,557	25	3,979	37,561
(%)	(56.4)	(43.6)	(100.0)			
60～	12,220	7,564	19,784	24	2,428	22,236
(%)	(61.8)	(38.2)	(100.0)			
70～	6,237	3,419	9,656	13	1,323	10,992
(%)	(64.6)	(35.4)	(100.0)			
80～	2,995	1,577	4,572	7	650	5,229
(%)	(65.5)	(34.5)	(100.0)			
90～	1,319	680	1,999	2	330	2,331
(%)	(66.0)	(34.0)	(100.0)			
100～	1,097	607	1,704	1	296	2,001
(%)	(64.4)	(35.6)	(100.0)			
合計	88,719	86,650	175,369	175	18,424	193,968
(%)	(50.6)	(49.4)	(100.0)			
記載なし	41,185	49,776	90,961	183	24,996	116,140
(%)	(45.3)	(54.7)	(100.0)			
総計	129,904	136,426	266,330	358	43,420	310,108
(%)	(48.8)	(51.2)	(100.0)			
平均	51.32	45.47	48.43	49.99	51.75	48.75
標準偏差	16.87	15.68	16.55	16.42	17.81	16.71

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査