

表 585 糖尿病の既往 透析前脈拍(拍/分)別 (透析患者全体)

透析前脈拍	既往なし	既往あり	合計	不明	記載なし	総計
40未満	64	102	166		19	185
(%)	(38.6)	(61.4)	(100.0)			
40～	1,123	1,046	2,169		212	2,381
(%)	(51.8)	(48.2)	(100.0)			
50～	9,729	7,861	17,590	26	1,896	19,512
(%)	(55.3)	(44.7)	(100.0)			
60～	29,539	25,251	54,790	58	5,933	60,781
(%)	(53.9)	(46.1)	(100.0)			
70～	33,889	32,584	66,473	75	6,724	73,272
(%)	(51.0)	(49.0)	(100.0)			
80～	19,662	22,890	42,552	66	4,196	46,814
(%)	(46.2)	(53.8)	(100.0)			
90～	8,290	10,581	18,871	24	1,695	20,590
(%)	(43.9)	(56.1)	(100.0)			
100～	2,631	3,256	5,887	10	496	6,393
(%)	(44.7)	(55.3)	(100.0)			
110～	707	691	1,398		130	1,528
(%)	(50.6)	(49.4)	(100.0)			
120～	339	259	598		57	655
(%)	(56.7)	(43.3)	(100.0)			
合計	105,973	104,521	210,494	259	21,358	232,111
(%)	(50.3)	(49.7)	(100.0)			
記載なし	23,931	31,905	55,836	99	22,062	77,997
(%)	(42.9)	(57.1)	(100.0)			
総計	129,904	136,426	266,330	358	43,420	310,108
(%)	(48.8)	(51.2)	(100.0)			
平均	73.93	75.59	74.75	75.72	74.05	74.69
標準偏差	12.77	12.90	12.86	12.38	12.59	12.84

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査