

表 601 糖尿病の既往 血流量(ml/分)別（施設血液透析患者 透析回数週3回）

血流量	既往なし	既往あり	合計	不明	記載なし	総計
100未満	16	25	41		3	44
(%)	(39.0)	(61.0)	(100.0)			
100～	316	361	677	1	61	739
(%)	(46.7)	(53.3)	(100.0)			
120～	926	1,027	1,953	10	185	2,148
(%)	(47.4)	(52.6)	(100.0)			
140～	7,699	8,635	16,334	25	1,614	17,973
(%)	(47.1)	(52.9)	(100.0)			
160～	2,868	2,582	5,450	3	448	5,901
(%)	(52.6)	(47.4)	(100.0)			
180～	13,486	14,207	27,693	30	2,762	30,485
(%)	(48.7)	(51.3)	(100.0)			
200～	44,067	47,147	91,214	65	9,211	100,490
(%)	(48.3)	(51.7)	(100.0)			
220～	13,473	13,219	26,692	22	2,832	29,546
(%)	(50.5)	(49.5)	(100.0)			
240～	12,275	12,245	24,520	29	2,511	27,060
(%)	(50.1)	(49.9)	(100.0)			
260～	1,135	1,130	2,265	2	266	2,533
(%)	(50.1)	(49.9)	(100.0)			
280～	1,045	1,065	2,110	4	223	2,337
(%)	(49.5)	(50.5)	(100.0)			
300～	1,852	1,899	3,751	5	509	4,265
(%)	(49.4)	(50.6)	(100.0)			
350～	205	198	403		26	429
(%)	(50.9)	(49.1)	(100.0)			
400～	353	318	671	2	45	718
(%)	(52.6)	(47.4)	(100.0)			
合計	99,716	104,058	203,774	198	20,696	224,668
(%)	(48.9)	(51.1)	(100.0)			
記載なし	742	1,818	2,560	32	1,711	4,303
(%)	(29.0)	(71.0)	(100.0)			
総計	100,458	105,876	206,334	230	22,407	228,971
(%)	(48.7)	(51.3)	(100.0)			
平均	205.35	204.36	204.85	201.57	205.86	204.94
標準偏差	36.74	36.23	36.49	44.64	35.64	36.42

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査