

表 606 糖尿病の既往 Body Mass Index別（施設血液透析患者 透析回数週3回）

Body Mass Index	既往なし	既往あり	合計	不明	記載なし	総計
14未満	744	419	1,163		148	1,311
(%)	(64.0)	(36.0)	(100.0)			
14～	4,400	2,435	6,835	14	757	7,606
(%)	(64.4)	(35.6)	(100.0)			
16～	13,589	8,490	22,079	30	2,371	24,480
(%)	(61.5)	(38.5)	(100.0)			
18～	22,354	16,959	39,313	34	3,809	43,156
(%)	(56.9)	(43.1)	(100.0)			
20～	22,618	20,972	43,590	33	4,057	47,680
(%)	(51.9)	(48.1)	(100.0)			
22～	15,228	18,712	33,940	22	2,848	36,810
(%)	(44.9)	(55.1)	(100.0)			
24～	8,066	12,340	20,406	15	1,492	21,913
(%)	(39.5)	(60.5)	(100.0)			
26～	3,713	7,264	10,977	6	705	11,688
(%)	(33.8)	(66.2)	(100.0)			
28～	1,716	3,940	5,656	8	327	5,991
(%)	(30.3)	(69.7)	(100.0)			
30～	1,363	3,538	4,901	1	262	5,164
(%)	(27.8)	(72.2)	(100.0)			
合計	93,791	95,069	188,860	163	16,776	205,799
(%)	(49.7)	(50.3)	(100.0)			
記載なし	6,667	10,807	17,474	67	5,631	23,172
(%)	(38.2)	(61.8)	(100.0)			
総計	100,458	105,876	206,334	230	22,407	228,971
(%)	(48.7)	(51.3)	(100.0)			
平均	20.81	22.21	21.51	20.71	20.91	21.46
標準偏差	3.40	3.80	3.67	3.66	3.43	3.66

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査