

表 607 糖尿病の既往 透析前BUN(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前BUN	既往なし	既往あり	合計	不明	記載なし	総計
20未満	183	251	434		45	479
(%)	(42.2)	(57.8)	(100.0)			
20～	915	1,355	2,270	5	205	2,480
(%)	(40.3)	(59.7)	(100.0)			
30～	3,869	5,617	9,486	5	867	10,358
(%)	(40.8)	(59.2)	(100.0)			
40～	10,901	13,721	24,622	27	2,162	26,811
(%)	(44.3)	(55.7)	(100.0)			
50～	18,971	19,751	38,722	25	3,912	42,659
(%)	(49.0)	(51.0)	(100.0)			
60～	20,652	17,826	38,478	28	4,137	42,643
(%)	(53.7)	(46.3)	(100.0)			
70～	14,668	10,803	25,471	23	2,828	28,322
(%)	(57.6)	(42.4)	(100.0)			
80～	7,015	4,782	11,797	13	1,402	13,212
(%)	(59.5)	(40.5)	(100.0)			
90～	2,584	1,679	4,263	4	536	4,803
(%)	(60.6)	(39.4)	(100.0)			
100～	728	480	1,208	2	138	1,348
(%)	(60.3)	(39.7)	(100.0)			
110～	202	151	353		41	394
(%)	(57.2)	(42.8)	(100.0)			
120～	106	78	184		20	204
(%)	(57.6)	(42.4)	(100.0)			
合計	80,794	76,494	157,288	132	16,293	173,713
(%)	(51.4)	(48.6)	(100.0)			
記載なし	1,044	1,796	2,840	26	1,630	4,496
(%)	(36.8)	(63.2)	(100.0)			
総計	81,838	78,290	160,128	158	17,923	178,209
(%)	(51.1)	(48.9)	(100.0)			
平均	62.52	59.00	60.81	61.01	62.19	60.94
標準偏差	15.52	15.54	15.63	16.67	15.69	15.64

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査