

表 611 糖尿病の既往 nPCR(g/kg/day)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

nPCR	既往なし	既往あり	合計	不明	記載なし	総計
0.4未満	140	202	342		27	369
(%)	(40.9)	(59.1)	(100.0)			
0.4～	2,950	4,786	7,736	8	674	8,418
(%)	(38.1)	(61.9)	(100.0)			
0.6～	20,328	26,547	46,875	41	4,040	50,956
(%)	(43.4)	(56.6)	(100.0)			
0.8～	34,055	29,704	63,759	40	6,712	70,511
(%)	(53.4)	(46.6)	(100.0)			
1.0～	15,677	9,491	25,168	27	3,053	28,248
(%)	(62.3)	(37.7)	(100.0)			
1.2～	2,874	1,453	4,327	7	582	4,916
(%)	(66.4)	(33.6)	(100.0)			
1.4～	323	160	483	1	71	555
(%)	(66.9)	(33.1)	(100.0)			
1.6～	43	26	69		10	79
(%)	(62.3)	(37.7)	(100.0)			
1.8～	18	5	23		2	25
(%)	(78.3)	(21.7)	(100.0)			
2.0～	15	22	37		1	38
(%)	(40.5)	(59.5)	(100.0)			
合計	76,423	72,396	148,819	124	15,172	164,115
(%)	(51.4)	(48.6)	(100.0)			
記載なし	5,415	5,894	11,309	34	2,751	14,094
(%)	(47.9)	(52.1)	(100.0)			
総計	81,838	78,290	160,128	158	17,923	178,209
(%)	(51.1)	(48.9)	(100.0)			
平均	0.88	0.83	0.86	0.87	0.88	0.86
標準偏差	0.18	0.17	0.18	0.20	0.18	0.18

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査