

表 618 糖尿病の既往 PTH値(pg/mL)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	645	744	1,389		110	1,499
(%)	(46.4)	(53.6)	(100.0)			
20～	851	884	1,735		113	1,848
(%)	(49.0)	(51.0)	(100.0)			
35～	727	753	1,480	1	132	1,613
(%)	(49.1)	(50.9)	(100.0)			
50～	1,110	1,166	2,276		183	2,459
(%)	(48.8)	(51.2)	(100.0)			
75～	980	982	1,962		186	2,148
(%)	(49.9)	(50.1)	(100.0)			
100～	797	836	1,633		154	1,787
(%)	(48.8)	(51.2)	(100.0)			
125～	527	519	1,046		141	1,187
(%)	(50.4)	(49.6)	(100.0)			
150～	14	19	33		4	37
(%)	(42.4)	(57.6)	(100.0)			
151～	443	396	839	1	94	934
(%)	(52.8)	(47.2)	(100.0)			
180～	312	303	615		51	666
(%)	(50.7)	(49.3)	(100.0)			
210～	205	223	428		44	472
(%)	(47.9)	(52.1)	(100.0)			
240～	279	196	475		51	526
(%)	(58.7)	(41.3)	(100.0)			
300～	105	77	182		23	205
(%)	(57.7)	(42.3)	(100.0)			
350～	64	58	122		13	135
(%)	(52.5)	(47.5)	(100.0)			
400～	83	52	135		17	152
(%)	(61.5)	(38.5)	(100.0)			
500～	97	56	153		13	166
(%)	(63.4)	(36.6)	(100.0)			
合計	7,239	7,264	14,503	2	1,329	15,834
(%)	(49.9)	(50.1)	(100.0)			
記載なし	112	192	304	1	133	438
(%)	(36.8)	(63.2)	(100.0)			
総計	7,351	7,456	14,807	3	1,462	16,272
(%)	(49.6)	(50.4)	(100.0)			
平均	110.45	99.42	104.93	104.00	114.11	105.70
標準偏差	122.57	95.47	109.97	89.10	100.67	109.24

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査