

表 619 糖尿病の既往 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前ヘモグロビン 濃度	既往なし	既往あり	合計	不明	記載なし	総計
5未満	143	104	247		14	261
(%)	(57.9)	(42.1)	(100.0)			
5～	139	201	340		22	362
(%)	(40.9)	(59.1)	(100.0)			
6～	373	482	855	5	82	942
(%)	(43.6)	(56.4)	(100.0)			
7～	1,541	1,712	3,253	7	318	3,578
(%)	(47.4)	(52.6)	(100.0)			
8～	5,050	5,672	10,722	15	989	11,726
(%)	(47.1)	(52.9)	(100.0)			
9～	16,275	17,469	33,744	36	3,258	37,038
(%)	(48.2)	(51.8)	(100.0)			
10～	33,951	34,521	68,472	62	6,683	75,217
(%)	(49.6)	(50.4)	(100.0)			
11～	27,708	28,205	55,913	44	5,522	61,479
(%)	(49.6)	(50.4)	(100.0)			
12～	10,045	10,408	20,453	19	1,898	22,370
(%)	(49.1)	(50.9)	(100.0)			
13～	2,401	2,657	5,058	2	477	5,537
(%)	(47.5)	(52.5)	(100.0)			
14～	838	936	1,774		159	1,933
(%)	(47.2)	(52.8)	(100.0)			
合計	98,464	102,367	200,831	190	19,422	220,443
(%)	(49.0)	(51.0)	(100.0)			
記載なし	1,994	3,509	5,503	40	2,985	8,528
(%)	(36.2)	(63.8)	(100.0)			
総計	100,458	105,876	206,334	230	22,407	228,971
(%)	(48.7)	(51.3)	(100.0)			
平均	10.70	10.68	10.69	10.35	10.69	10.69
標準偏差	1.26	1.29	1.28	1.38	1.25	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査