

表 621 糖尿病の既往 HDL-C濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

HDL-C濃度	既往なし	既往あり	合計	不明	記載なし	総計
20未満	223	402	625		46	671
(%)	(35.7)	(64.3)	(100.0)			
20～	3,862	8,202	12,064	11	771	12,846
(%)	(32.0)	(68.0)	(100.0)			
30～	14,018	20,115	34,133	30	2,725	36,888
(%)	(41.1)	(58.9)	(100.0)			
40～	18,357	18,963	37,320	38	3,371	40,729
(%)	(49.2)	(50.8)	(100.0)			
50～	14,435	11,799	26,234	20	2,826	29,080
(%)	(55.0)	(45.0)	(100.0)			
60～	9,040	6,036	15,076	18	1,698	16,792
(%)	(60.0)	(40.0)	(100.0)			
70～	4,580	2,708	7,288	10	903	8,201
(%)	(62.8)	(37.2)	(100.0)			
80～	2,137	1,236	3,373	5	456	3,834
(%)	(63.4)	(36.6)	(100.0)			
90～	949	542	1,491	1	241	1,733
(%)	(63.6)	(36.4)	(100.0)			
100～	759	482	1,241		217	1,458
(%)	(61.2)	(38.8)	(100.0)			
合計	68,360	70,485	138,845	133	13,254	152,232
(%)	(49.2)	(50.8)	(100.0)			
記載なし	32,098	35,391	67,489	97	9,153	76,739
(%)	(47.6)	(52.4)	(100.0)			
総計	100,458	105,876	206,334	230	22,407	228,971
(%)	(48.7)	(51.3)	(100.0)			
平均	50.72	45.27	47.95	49.13	51.43	48.25
標準偏差	16.68	15.60	16.37	15.52	17.95	16.54

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査