

表 625 糖尿病の既往 透析前脈拍(拍/分)別（施設血液透析患者 透析回数週3回）

透析前脈拍	既往なし	既往あり	合計	不明	記載なし	総計
40未満	52	89	141		14	155
(%)	(36.9)	(63.1)	(100.0)			
40～	900	870	1,770		159	1,929
(%)	(50.8)	(49.2)	(100.0)			
50～	7,779	6,586	14,365	22	1,386	15,773
(%)	(54.2)	(45.8)	(100.0)			
60～	23,506	20,906	44,412	42	4,287	48,741
(%)	(52.9)	(47.1)	(100.0)			
70～	26,463	26,790	53,253	53	4,878	58,184
(%)	(49.7)	(50.3)	(100.0)			
80～	15,037	18,655	33,692	44	2,987	36,723
(%)	(44.6)	(55.4)	(100.0)			
90～	6,261	8,443	14,704	13	1,201	15,918
(%)	(42.6)	(57.4)	(100.0)			
100～	1,966	2,570	4,536	4	364	4,904
(%)	(43.3)	(56.7)	(100.0)			
110～	511	555	1,066		91	1,157
(%)	(47.9)	(52.1)	(100.0)			
120～	259	201	460		38	498
(%)	(56.3)	(43.7)	(100.0)			
合計	82,734	85,665	168,399	178	15,405	183,982
(%)	(49.1)	(50.9)	(100.0)			
記載なし	17,724	20,211	37,935	52	7,002	44,989
(%)	(46.7)	(53.3)	(100.0)			
総計	100,458	105,876	206,334	230	22,407	228,971
(%)	(48.7)	(51.3)	(100.0)			
平均	73.67	75.40	74.55	74.11	73.96	74.50
標準偏差	12.71	12.86	12.82	12.12	12.59	12.80

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査