

表 642 糖尿病の既往 身長(cm)別（血液透析濾過患者 透析回数週3回）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	19	15	34		4	38
(%)	(55.9)	(44.1)	(100.0)			
130～	38	14	52	1	17	70
(%)	(73.1)	(26.9)	(100.0)			
135～	172	81	253	1	35	289
(%)	(68.0)	(32.0)	(100.0)			
140～	607	420	1,027	3	133	1,163
(%)	(59.1)	(40.9)	(100.0)			
145～	1,614	1,040	2,654	7	334	2,995
(%)	(60.8)	(39.2)	(100.0)			
150～	2,953	1,693	4,646	6	560	5,212
(%)	(63.6)	(36.4)	(100.0)			
155～	3,285	2,174	5,459	11	609	6,079
(%)	(60.2)	(39.8)	(100.0)			
160～	3,548	2,971	6,519	6	652	7,177
(%)	(54.4)	(45.6)	(100.0)			
165～	3,267	2,857	6,124	4	594	6,722
(%)	(53.3)	(46.7)	(100.0)			
170～	2,291	2,089	4,380	1	404	4,785
(%)	(52.3)	(47.7)	(100.0)			
175～	864	764	1,628	2	156	1,786
(%)	(53.1)	(46.9)	(100.0)			
180～	235	218	453		40	493
(%)	(51.9)	(48.1)	(100.0)			
185～	32	36	68		9	77
(%)	(47.1)	(52.9)	(100.0)			
190～	2	7	9			9
(%)	(22.2)	(77.8)	(100.0)			
合計	18,927	14,379	33,306	42	3,547	36,895
(%)	(56.8)	(43.2)	(100.0)			
記載なし	758	1,288	2,046	8	997	3,051
(%)	(37.0)	(63.0)	(100.0)			
総計	19,685	15,667	35,352	50	4,544	39,946
(%)	(55.7)	(44.3)	(100.0)			
平均	160.23	161.66	160.85	155.19	159.74	160.74
標準偏差	9.54	9.36	9.49	10.00	9.88	9.54

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査