

表 643 糖尿病の既往 透析後体重(kg)別（血液透析濾過患者 透析回数週3回）

透析後体重	既往なし	既往あり	合計	不明	記載なし	総計
20未満	6	3	9		3	12
(%)	(66.7)	(33.3)	(100.0)			
20～	48	15	63		13	76
(%)	(76.2)	(23.8)	(100.0)			
30～	1,664	629	2,293	10	379	2,682
(%)	(72.6)	(27.4)	(100.0)			
40～	5,837	2,919	8,756	16	1,233	10,005
(%)	(66.7)	(33.3)	(100.0)			
50～	6,294	4,832	11,126	10	1,354	12,490
(%)	(56.6)	(43.4)	(100.0)			
60～	3,754	3,825	7,579	3	783	8,365
(%)	(49.5)	(50.5)	(100.0)			
70～	1,333	1,820	3,153	2	268	3,423
(%)	(42.3)	(57.7)	(100.0)			
80～	399	802	1,201	1	71	1,273
(%)	(33.2)	(66.8)	(100.0)			
90～	117	312	429		26	455
(%)	(27.3)	(72.7)	(100.0)			
100～	33	97	130		12	142
(%)	(25.4)	(74.6)	(100.0)			
110～	17	41	58		3	61
(%)	(29.3)	(70.7)	(100.0)			
120～	13	22	35		2	37
(%)	(37.1)	(62.9)	(100.0)			
合計	19,515	15,317	34,832	42	4,147	39,021
(%)	(56.0)	(44.0)	(100.0)			
記載なし	170	350	520	8	397	925
(%)	(32.7)	(67.3)	(100.0)			
総計	19,685	15,667	35,352	50	4,544	39,946
(%)	(55.7)	(44.3)	(100.0)			
平均	54.49	60.04	56.93	48.63	54.15	56.63
標準偏差	12.00	13.80	13.11	11.08	12.03	13.03

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査