

表 646 糖尿病の既往 Body Mass Index別（血液透析濾過患者 透析回数週3回）

Body Mass Index	既往なし	既往あり	合計	不明	記載なし	総計
14未満	98	34	132		17	149
(%)	(74.2)	(25.8)	(100.0)			
14～	644	261	905	1	140	1,046
(%)	(71.2)	(28.8)	(100.0)			
16～	2,535	991	3,526	14	458	3,998
(%)	(71.9)	(28.1)	(100.0)			
18～	4,456	2,318	6,774	8	832	7,614
(%)	(65.8)	(34.2)	(100.0)			
20～	4,610	3,009	7,619	4	839	8,462
(%)	(60.5)	(39.5)	(100.0)			
22～	3,220	2,784	6,004	6	561	6,571
(%)	(53.6)	(46.4)	(100.0)			
24～	1,674	2,082	3,756	5	327	4,088
(%)	(44.6)	(55.4)	(100.0)			
26～	830	1,216	2,046		151	2,197
(%)	(40.6)	(59.4)	(100.0)			
28～	380	731	1,111	2	64	1,177
(%)	(34.2)	(65.8)	(100.0)			
30～	317	662	979		38	1,017
(%)	(32.4)	(67.6)	(100.0)			
合計	18,764	14,088	32,852	40	3,427	36,319
(%)	(57.1)	(42.9)	(100.0)			
記載なし	921	1,579	2,500	10	1,117	3,627
(%)	(36.8)	(63.2)	(100.0)			
総計	19,685	15,667	35,352	50	4,544	39,946
(%)	(55.7)	(44.3)	(100.0)			
平均	21.05	22.68	21.75	20.35	20.93	21.67
標準偏差	3.38	3.84	3.67	3.53	3.30	3.65

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査