

表 647 糖尿病の既往 透析前BUN(mg/dL)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

透析前BUN	既往なし	既往あり	合計	不明	記載なし	総計
20未満	16	17	33		6	39
(%)	(48.5)	(51.5)	(100.0)			
20～	121	175	296		13	309
(%)	(40.9)	(59.1)	(100.0)			
30～	658	846	1,504	2	147	1,653
(%)	(43.8)	(56.3)	(100.0)			
40～	2,146	2,159	4,305	4	476	4,785
(%)	(49.8)	(50.2)	(100.0)			
50～	4,131	3,305	7,436	9	873	8,318
(%)	(55.6)	(44.4)	(100.0)			
60～	4,835	2,983	7,818	13	973	8,804
(%)	(61.8)	(38.2)	(100.0)			
70～	3,477	1,883	5,360	6	720	6,086
(%)	(64.9)	(35.1)	(100.0)			
80～	1,603	896	2,499	2	330	2,831
(%)	(64.1)	(35.9)	(100.0)			
90～	616	303	919	3	129	1,051
(%)	(67.0)	(33.0)	(100.0)			
100～	166	88	254		25	279
(%)	(65.4)	(34.6)	(100.0)			
110～	43	19	62		12	74
(%)	(69.4)	(30.6)	(100.0)			
120～	12	6	18		2	20
(%)	(66.7)	(33.3)	(100.0)			
合計	17,824	12,680	30,504	39	3,706	34,249
(%)	(58.4)	(41.6)	(100.0)			
記載なし	81	255	336	8	378	722
(%)	(24.1)	(75.9)	(100.0)			
総計	17,905	12,935	30,840	47	4,084	34,971
(%)	(58.1)	(41.9)	(100.0)			
平均	63.66	59.87	62.09	63.69	63.42	62.23
標準偏差	14.81	15.25	15.11	14.03	14.83	15.08

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査