

表 656 糖尿病の既往 透析前リン濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

透析前リン濃度	既往なし	既往あり	合計	不明	記載なし	総計
3未満	541	485	1,026	1	106	1,133
(%)	(52.7)	(47.3)	(100.0)			
3.0～	776	644	1,420	1	153	1,574
(%)	(54.6)	(45.4)	(100.0)			
3.5～	1,331	1,234	2,565	2	282	2,849
(%)	(51.9)	(48.1)	(100.0)			
4.0～	2,267	1,832	4,099	5	461	4,565
(%)	(55.3)	(44.7)	(100.0)			
4.5～	2,837	2,256	5,093	6	613	5,712
(%)	(55.7)	(44.3)	(100.0)			
5.0～	3,184	2,276	5,460	9	665	6,134
(%)	(58.3)	(41.7)	(100.0)			
5.5～	2,759	1,972	4,731	7	561	5,299
(%)	(58.3)	(41.7)	(100.0)			
6.0～	516	327	843	1	106	950
(%)	(61.2)	(38.8)	(100.0)			
6.1～	1,630	1,267	2,897	1	346	3,244
(%)	(56.3)	(43.7)	(100.0)			
6.5～	1,400	1,101	2,501	2	303	2,806
(%)	(56.0)	(44.0)	(100.0)			
7.0～	884	721	1,605		170	1,775
(%)	(55.1)	(44.9)	(100.0)			
7.5～	544	455	999		105	1,104
(%)	(54.5)	(45.5)	(100.0)			
8.0～	341	255	596	1	54	651
(%)	(57.2)	(42.8)	(100.0)			
8.5～	210	173	383		40	423
(%)	(54.8)	(45.2)	(100.0)			
9.0～	352	273	625	1	56	682
(%)	(56.3)	(43.7)	(100.0)			
合計	19,572	15,271	34,843	37	4,021	38,901
(%)	(56.2)	(43.8)	(100.0)			
記載なし	113	396	509	13	523	1,045
(%)	(22.2)	(77.8)	(100.0)			
総計	19,685	15,667	35,352	50	4,544	39,946
(%)	(55.7)	(44.3)	(100.0)			
平均	5.38	5.33	5.36	5.16	5.34	5.36
標準偏差	1.41	1.45	1.43	1.31	1.37	1.43

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査