

表 658 糖尿病の既往 PTH値(pg/mL)別（血液透析濾過患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	124	98	222		27	249
(%)	(55.9)	(44.1)	(100.0)			
20～	139	134	273		28	301
(%)	(50.9)	(49.1)	(100.0)			
35～	141	105	246		28	274
(%)	(57.3)	(42.7)	(100.0)			
50～	229	167	396		48	444
(%)	(57.8)	(42.2)	(100.0)			
75～	201	152	353		38	391
(%)	(56.9)	(43.1)	(100.0)			
100～	166	126	292		39	331
(%)	(56.8)	(43.2)	(100.0)			
125～	124	82	206		28	234
(%)	(60.2)	(39.8)	(100.0)			
150～	4		4			4
(%)	(100.0)		(100.0)			
151～	117	78	195		20	215
(%)	(60.0)	(40.0)	(100.0)			
180～	87	41	128		9	137
(%)	(68.0)	(32.0)	(100.0)			
210～	56	33	89		9	98
(%)	(62.9)	(37.1)	(100.0)			
240～	50	31	81		7	88
(%)	(61.7)	(38.3)	(100.0)			
300～	21	16	37		5	42
(%)	(56.8)	(43.2)	(100.0)			
350～	20	7	27		2	29
(%)	(74.1)	(25.9)	(100.0)			
400～	14	6	20		4	24
(%)	(70.0)	(30.0)	(100.0)			
500～	33	9	42		3	45
(%)	(78.6)	(21.4)	(100.0)			
合計	1,526	1,085	2,611		295	2,906
(%)	(58.4)	(41.6)	(100.0)			
記載なし	7	4	11		20	31
(%)	(63.6)	(36.4)	(100.0)			
総計	1,533	1,089	2,622		315	2,937
(%)	(58.5)	(41.5)	(100.0)			
平均	120.21	101.10	112.27		107.56	111.79
標準偏差	120.96	88.08	108.90		100.96	108.12

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査