

表 659 糖尿病の既往 透析前ヘモグロビン濃度(g/dL)別（血液透析濾過患者 透析回数週3回）

透析前ヘモグロビン 濃度	既往なし	既往あり	合計	不明	記載なし	総計
5未満	15	11	26			26
(%)	(57.7)	(42.3)	(100.0)			
5～	17	12	29		2	31
(%)	(58.6)	(41.4)	(100.0)			
6～	62	42	104		10	114
(%)	(59.6)	(40.4)	(100.0)			
7～	218	184	402		43	445
(%)	(54.2)	(45.8)	(100.0)			
8～	790	681	1,471	2	164	1,637
(%)	(53.7)	(46.3)	(100.0)			
9～	2,826	2,379	5,205	6	639	5,850
(%)	(54.3)	(45.7)	(100.0)			
10～	6,490	5,055	11,545	18	1,384	12,947
(%)	(56.2)	(43.8)	(100.0)			
11～	5,933	4,431	10,364	8	1,205	11,577
(%)	(57.2)	(42.8)	(100.0)			
12～	2,296	1,672	3,968	2	412	4,382
(%)	(57.9)	(42.1)	(100.0)			
13～	578	496	1,074	1	110	1,185
(%)	(53.8)	(46.2)	(100.0)			
14～	254	229	483		40	523
(%)	(52.6)	(47.4)	(100.0)			
合計	19,479	15,192	34,671	37	4,009	38,717
(%)	(56.2)	(43.8)	(100.0)			
記載なし	206	475	681	13	535	1,229
(%)	(30.2)	(69.8)	(100.0)			
総計	19,685	15,667	35,352	50	4,544	39,946
(%)	(55.7)	(44.3)	(100.0)			
平均	10.86	10.83	10.84	10.50	10.80	10.84
標準偏差	1.25	1.29	1.27	0.99	1.20	1.26

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査