

表 661 糖尿病の既往 HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

HDL-C濃度	既往なし	既往あり	合計	不明	記載なし	総計
20未満	22	54	76		5	81
(%)	(28.9)	(71.1)	(100.0)			
20～	575	1,147	1,722		152	1,874
(%)	(33.4)	(66.6)	(100.0)			
30～	2,448	3,166	5,614	3	514	6,131
(%)	(43.6)	(56.4)	(100.0)			
40～	3,618	3,012	6,630	3	727	7,360
(%)	(54.6)	(45.4)	(100.0)			
50～	3,256	2,023	5,279	1	691	5,971
(%)	(61.7)	(38.3)	(100.0)			
60～	2,356	1,057	3,413	2	442	3,857
(%)	(69.0)	(31.0)	(100.0)			
70～	1,223	518	1,741	1	281	2,023
(%)	(70.2)	(29.8)	(100.0)			
80～	639	243	882		131	1,013
(%)	(72.4)	(27.6)	(100.0)			
90～	274	97	371	1	59	431
(%)	(73.9)	(26.1)	(100.0)			
100～	243	83	326		57	383
(%)	(74.5)	(25.5)	(100.0)			
合計	14,654	11,400	26,054	11	3,059	29,124
(%)	(56.2)	(43.8)	(100.0)			
記載なし	5,031	4,267	9,298	39	1,485	10,822
(%)	(54.1)	(45.9)	(100.0)			
総計	19,685	15,667	35,352	50	4,544	39,946
(%)	(55.7)	(44.3)	(100.0)			
平均	53.75	46.35	50.52	55.82	53.72	50.85
標準偏差	17.28	15.86	17.07	18.02	17.86	17.18

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査