

表 662 糖尿病の既往 Non-HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

Non-HDL-C濃度	既往なし	既往あり	合計	不明	記載なし	総計
50未満	221	236	457		72	529
(%)	(48.4)	(51.6)	(100.0)			
50～	1,054	1,079	2,133	1	269	2,403
(%)	(49.4)	(50.6)	(100.0)			
70～	2,495	2,095	4,590	5	543	5,138
(%)	(54.4)	(45.6)	(100.0)			
90～	3,203	2,390	5,593	2	670	6,265
(%)	(57.3)	(42.7)	(100.0)			
110～	2,650	1,931	4,581	2	579	5,162
(%)	(57.8)	(42.2)	(100.0)			
130～	1,695	1,113	2,808		341	3,149
(%)	(60.4)	(39.6)	(100.0)			
150～	806	591	1,397		200	1,597
(%)	(57.7)	(42.3)	(100.0)			
170～	350	267	617		65	682
(%)	(56.7)	(43.3)	(100.0)			
190～	127	109	236		26	262
(%)	(53.8)	(46.2)	(100.0)			
210～	61	57	118		23	141
(%)	(51.7)	(48.3)	(100.0)			
合計	12,662	9,868	22,530	10	2,788	25,328
(%)	(56.2)	(43.8)	(100.0)			
記載なし	7,023	5,799	12,822	40	1,756	14,618
(%)	(54.8)	(45.2)	(100.0)			
総計	19,685	15,667	35,352	50	4,544	39,946
(%)	(55.7)	(44.3)	(100.0)			
平均	108.41	105.39	107.08	88.70	107.46	107.12
標準偏差	32.75	33.88	33.29	17.57	34.08	33.37

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査