

表 664 糖尿病の既往 透析前拡張期血圧(mmHg)別（血液透析濾過患者 透析回数週3回）

透析前拡張 期血圧	既往なし	既往あり	合計	不明	記載なし	総計
20未満	4	2	6		1	7
(%)	(66.7)	(33.3)	(100.0)			
20～	59	42	101		20	121
(%)	(58.4)	(41.6)	(100.0)			
40～	1,348	1,066	2,414	4	262	2,680
(%)	(55.8)	(44.2)	(100.0)			
60～	7,235	6,184	13,419	15	1,538	14,972
(%)	(53.9)	(46.1)	(100.0)			
80～	4,633	3,392	8,025	6	972	9,003
(%)	(57.7)	(42.3)	(100.0)			
90～	2,759	1,998	4,757	8	552	5,317
(%)	(58.0)	(42.0)	(100.0)			
100～	1,358	1,072	2,430	3	268	2,701
(%)	(55.9)	(44.1)	(100.0)			
120～	82	70	152		17	169
(%)	(53.9)	(46.1)	(100.0)			
140～	5	7	12			12
(%)	(41.7)	(58.3)	(100.0)			
合計	17,483	13,833	31,316	36	3,630	34,982
(%)	(55.8)	(44.2)	(100.0)			
記載なし	2,202	1,834	4,036	14	914	4,964
(%)	(54.6)	(45.4)	(100.0)			
総計	19,685	15,667	35,352	50	4,544	39,946
(%)	(55.7)	(44.3)	(100.0)			
平均	79.38	78.93	79.18	78.81	79.08	79.17
標準偏差	14.71	14.74	14.73	16.21	14.47	14.70

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査