

表 665 糖尿病の既往 透析前脈拍(拍/分)別（血液透析濾過患者 透析回数週3回）

透析前脈拍	既往なし	既往あり	合計	不明	記載なし	総計
40未満	6	8	14		3	17
(%)	(42.9)	(57.1)	(100.0)			
40～	151	118	269		26	295
(%)	(56.1)	(43.9)	(100.0)			
50～	1,438	884	2,322	3	307	2,632
(%)	(61.9)	(38.1)	(100.0)			
60～	4,444	3,064	7,508	5	977	8,490
(%)	(59.2)	(40.8)	(100.0)			
70～	5,381	4,050	9,431	12	1,102	10,545
(%)	(57.1)	(42.9)	(100.0)			
80～	3,289	3,036	6,325	10	728	7,063
(%)	(52.0)	(48.0)	(100.0)			
90～	1,410	1,564	2,974	5	314	3,293
(%)	(47.4)	(52.6)	(100.0)			
100～	458	495	953	1	88	1,042
(%)	(48.1)	(51.9)	(100.0)			
110～	131	108	239		22	261
(%)	(54.8)	(45.2)	(100.0)			
120～	50	46	96		10	106
(%)	(52.1)	(47.9)	(100.0)			
合計	16,758	13,373	30,131	36	3,577	33,744
(%)	(55.6)	(44.4)	(100.0)			
記載なし	2,927	2,294	5,221	14	967	6,202
(%)	(56.1)	(43.9)	(100.0)			
総計	19,685	15,667	35,352	50	4,544	39,946
(%)	(55.7)	(44.3)	(100.0)			
平均	74.61	76.69	75.53	78.61	74.51	75.43
標準偏差	12.78	13.09	12.96	11.54	12.65	12.93

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査