

表 679 糖尿病の既往 身長(cm)別（腹膜透析患者）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	53	7	60		2	62
(%)	(88.3)	(11.7)	(100.0)			
130～	10	1	11			11
(%)	(90.9)	(9.1)	(100.0)			
135～	41	17	58		4	62
(%)	(70.7)	(29.3)	(100.0)			
140～	122	83	205	1	21	227
(%)	(59.5)	(40.5)	(100.0)			
145～	264	154	418	1	51	470
(%)	(63.2)	(36.8)	(100.0)			
150～	441	314	755	4	108	867
(%)	(58.4)	(41.6)	(100.0)			
155～	601	353	954	2	93	1,049
(%)	(63.0)	(37.0)	(100.0)			
160～	632	553	1,185	5	140	1,330
(%)	(53.3)	(46.7)	(100.0)			
165～	551	529	1,080	1	131	1,212
(%)	(51.0)	(49.0)	(100.0)			
170～	413	400	813	2	89	904
(%)	(50.8)	(49.2)	(100.0)			
175～	167	165	332	1	33	366
(%)	(50.3)	(49.7)	(100.0)			
180～	43	39	82		5	87
(%)	(52.4)	(47.6)	(100.0)			
185～	6	5	11	1	1	13
(%)	(54.5)	(45.5)	(100.0)			
190～		3	3		1	4
(%)		(100.0)	(100.0)			
合計	3,344	2,623	5,967	18	679	6,664
(%)	(56.0)	(44.0)	(100.0)			
記載なし	329	629	958	8	1,311	2,277
(%)	(34.3)	(65.7)	(100.0)			
総計	3,673	3,252	6,925	26	1,990	8,941
(%)	(53.0)	(47.0)	(100.0)			
平均	159.43	161.97	160.54	160.83	160.67	160.56
標準偏差	12.78	10.10	11.74	11.17	10.20	11.59

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査