

表 680 糖尿病の既往 体重(kg)別（腹膜透析患者）

体重	既往なし	既往あり	合計	不明	記載なし	総計
20未満	38		38			38
(%)	(100.0)		(100.0)			
20～	22		22		1	23
(%)	(100.0)		(100.0)			
30～	129	32	161	2	9	172
(%)	(80.1)	(19.9)	(100.0)			
40～	610	228	838	5	61	904
(%)	(72.8)	(27.2)	(100.0)			
50～	904	563	1,467	5	119	1,591
(%)	(61.6)	(38.4)	(100.0)			
60～	823	689	1,512	4	100	1,616
(%)	(54.4)	(45.6)	(100.0)			
70～	361	406	767	2	42	811
(%)	(47.1)	(52.9)	(100.0)			
80～	128	150	278	4	16	298
(%)	(46.0)	(54.0)	(100.0)			
90～	24	77	101		3	104
(%)	(23.8)	(76.2)	(100.0)			
100～	6	14	20			20
(%)	(30.0)	(70.0)	(100.0)			
110～	2	7	9		1	10
(%)	(22.2)	(77.8)	(100.0)			
120～	3	6	9			9
(%)	(33.3)	(66.7)	(100.0)			
合計	3,050	2,172	5,222	22	352	5,596
(%)	(58.4)	(41.6)	(100.0)			
記載なし	623	1,080	1,703	4	1,638	3,345
(%)	(36.6)	(63.4)	(100.0)			
総計	3,673	3,252	6,925	26	1,990	8,941
(%)	(53.0)	(47.0)	(100.0)			
平均	58.15	64.55	60.81	60.49	59.71	60.74
標準偏差	13.77	13.51	14.02	15.27	11.82	13.90

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査