

表 682 糖尿病の既往 BUN(mg/dL)別（腹膜透析患者）

BUN	既往なし	既往あり	合計	不明	記載なし	総計
20未満	14	11	25		4	29
(%)	(56.0)	(44.0)	(100.0)			
20～	77	101	178	1	4	183
(%)	(43.3)	(56.7)	(100.0)			
30～	294	329	623	2	47	672
(%)	(47.2)	(52.8)	(100.0)			
40～	718	573	1,291	5	96	1,392
(%)	(55.6)	(44.4)	(100.0)			
50～	874	609	1,483	5	125	1,613
(%)	(58.9)	(41.1)	(100.0)			
60～	664	389	1,053	5	92	1,150
(%)	(63.1)	(36.9)	(100.0)			
70～	335	171	506	1	42	549
(%)	(66.2)	(33.8)	(100.0)			
80～	144	79	223	2	20	245
(%)	(64.6)	(35.4)	(100.0)			
90～	51	30	81		8	89
(%)	(63.0)	(37.0)	(100.0)			
100～	19	18	37		3	40
(%)	(51.4)	(48.6)	(100.0)			
110～	5	2	7			7
(%)	(71.4)	(28.6)	(100.0)			
120～	5	1	6		1	7
(%)	(83.3)	(16.7)	(100.0)			
合計	3,200	2,313	5,513	21	442	5,976
(%)	(58.0)	(42.0)	(100.0)			
記載なし	473	939	1,412	5	1,548	2,965
(%)	(33.5)	(66.5)	(100.0)			
総計	3,673	3,252	6,925	26	1,990	8,941
(%)	(53.0)	(47.0)	(100.0)			
平均	56.23	52.80	54.79	54.62	56.30	54.90
標準偏差	15.94	15.51	15.85	16.34	15.40	15.82

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査