

表 686 糖尿病の既往 補正カルシウム濃度(mg/dL)別（腹膜透析患者）

補正 カルシウム濃度	既往なし	既往あり	合計	不明	記載なし	総計
7.0未満	15	12	27		2	29
(%)	(55.6)	(44.4)	(100.0)			
7.0～	18	15	33		3	36
(%)	(54.5)	(45.5)	(100.0)			
7.5～	67	51	118		9	127
(%)	(56.8)	(43.2)	(100.0)			
8.0～	125	133	258		28	286
(%)	(48.4)	(51.6)	(100.0)			
8.4～	44	40	84		2	86
(%)	(52.4)	(47.6)	(100.0)			
8.5～	466	376	842	5	59	906
(%)	(55.3)	(44.7)	(100.0)			
9.0～	862	638	1,500	5	97	1,602
(%)	(57.5)	(42.5)	(100.0)			
9.5～	894	609	1,503	6	120	1,629
(%)	(59.5)	(40.5)	(100.0)			
10.0～	108	70	178	1	12	191
(%)	(60.7)	(39.3)	(100.0)			
10.1～	311	192	503	2	52	557
(%)	(61.8)	(38.2)	(100.0)			
10.5～	160	103	263		33	296
(%)	(60.8)	(39.2)	(100.0)			
11.0～	53	49	102	1	8	111
(%)	(52.0)	(48.0)	(100.0)			
11.5～	17	10	27		2	29
(%)	(63.0)	(37.0)	(100.0)			
12.0～	19	5	24		3	27
(%)	(79.2)	(20.8)	(100.0)			
合計	3,159	2,303	5,462	20	430	5,912
(%)	(57.8)	(42.2)	(100.0)			
記載なし	514	949	1,463	6	1,560	3,029
(%)	(35.1)	(64.9)	(100.0)			
総計	3,673	3,252	6,925	26	1,990	8,941
(%)	(53.0)	(47.0)	(100.0)			
平均	9.43	9.36	9.40	9.52	9.49	9.41
標準偏差	0.79	0.78	0.79	0.63	0.87	0.80

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査