

表 687 糖尿病の既往 リン濃度(mg/dL)別（腹膜透析患者）

リン濃度	既往なし	既往あり	合計	不明	記載なし	総計
3未満	59	54	113		13	126
(%)	(52.2)	(47.8)	(100.0)			
3.0～	127	122	249		21	270
(%)	(51.0)	(49.0)	(100.0)			
3.5～	275	212	487	2	36	525
(%)	(56.5)	(43.5)	(100.0)			
4.0～	405	315	720	3	47	770
(%)	(56.3)	(43.8)	(100.0)			
4.5～	521	345	866	3	52	921
(%)	(60.2)	(39.8)	(100.0)			
5.0～	507	369	876	3	69	948
(%)	(57.9)	(42.1)	(100.0)			
5.5～	459	318	777		62	839
(%)	(59.1)	(40.9)	(100.0)			
6.0～	73	55	128		10	138
(%)	(57.0)	(43.0)	(100.0)			
6.1～	249	188	437	4	36	477
(%)	(57.0)	(43.0)	(100.0)			
6.5～	224	130	354	3	42	399
(%)	(63.3)	(36.7)	(100.0)			
7.0～	119	73	192		20	212
(%)	(62.0)	(38.0)	(100.0)			
7.5～	77	60	137	1	11	149
(%)	(56.2)	(43.8)	(100.0)			
8.0～	37	28	65		6	71
(%)	(56.9)	(43.1)	(100.0)			
8.5～	23	13	36		2	38
(%)	(63.9)	(36.1)	(100.0)			
9.0～	45	33	78	1	10	89
(%)	(57.7)	(42.3)	(100.0)			
合計	3,200	2,315	5,515	20	437	5,972
(%)	(58.0)	(42.0)	(100.0)			
記載なし	473	937	1,410	6	1,553	2,969
(%)	(33.5)	(66.5)	(100.0)			
総計	3,673	3,252	6,925	26	1,990	8,941
(%)	(53.0)	(47.0)	(100.0)			
平均	5.27	5.19	5.24	5.68	5.34	5.25
標準偏差	1.33	1.36	1.34	1.59	1.44	1.35

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査