

表 689 糖尿病の既往 PTH値(pg/mL)別（腹膜透析患者）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	19	21	40			40
(%)	(47.5)	(52.5)	(100.0)			
20～	24	21	45			45
(%)	(53.3)	(46.7)	(100.0)			
35～	30	24	54		1	55
(%)	(55.6)	(44.4)	(100.0)			
50～	43	33	76		1	77
(%)	(56.6)	(43.4)	(100.0)			
75～	46	23	69			69
(%)	(66.7)	(33.3)	(100.0)			
100～	39	32	71			71
(%)	(54.9)	(45.1)	(100.0)			
125～	30	15	45			45
(%)	(66.7)	(33.3)	(100.0)			
150～	1		1			1
(%)	(100.0)		(100.0)			
151～	23	20	43			43
(%)	(53.5)	(46.5)	(100.0)			
180～	18	14	32		2	34
(%)	(56.3)	(43.8)	(100.0)			
210～	14	13	27			27
(%)	(51.9)	(48.1)	(100.0)			
240～	14	9	23			23
(%)	(60.9)	(39.1)	(100.0)			
300～	15	4	19			19
(%)	(78.9)	(21.1)	(100.0)			
350～	6	2	8			8
(%)	(75.0)	(25.0)	(100.0)			
400～	5	6	11			11
(%)	(45.5)	(54.5)	(100.0)			
500～	8	3	11			11
(%)	(72.7)	(27.3)	(100.0)			
合計	335	240	575		4	579
(%)	(58.3)	(41.7)	(100.0)			
記載なし	64	53	117		23	140
(%)	(54.7)	(45.3)	(100.0)			
総計	399	293	692		27	719
(%)	(57.7)	(42.3)	(100.0)			
平均	140.16	122.17	132.65		119.25	132.56
標準偏差	142.17	109.45	129.72		76.31	129.39

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査