

表 690 糖尿病の既往 ヘモグロビン濃度(g/dL)別（腹膜透析患者）

| ヘモグロビン<br>濃度 | 既往なし   | 既往あり   | 合計      | 不明    | 記載なし  | 総計    |
|--------------|--------|--------|---------|-------|-------|-------|
| 5未満          | 3      | 4      | 7       |       |       | 7     |
| (%)          | (42.9) | (57.1) | (100.0) |       |       |       |
| 5～           | 4      | 10     | 14      |       |       | 14    |
| (%)          | (28.6) | (71.4) | (100.0) |       |       |       |
| 6～           | 12     | 13     | 25      |       | 3     | 28    |
| (%)          | (48.0) | (52.0) | (100.0) |       |       |       |
| 7～           | 59     | 55     | 114     |       | 10    | 124   |
| (%)          | (51.8) | (48.2) | (100.0) |       |       |       |
| 8～           | 167    | 145    | 312     | 3     | 29    | 344   |
| (%)          | (53.5) | (46.5) | (100.0) |       |       |       |
| 9～           | 474    | 357    | 831     | 7     | 65    | 903   |
| (%)          | (57.0) | (43.0) | (100.0) |       |       |       |
| 10～          | 934    | 643    | 1,577   | 3     | 126   | 1,706 |
| (%)          | (59.2) | (40.8) | (100.0) |       |       |       |
| 11～          | 954    | 632    | 1,586   | 6     | 117   | 1,709 |
| (%)          | (60.2) | (39.8) | (100.0) |       |       |       |
| 12～          | 413    | 320    | 733     | 2     | 61    | 796   |
| (%)          | (56.3) | (43.7) | (100.0) |       |       |       |
| 13～          | 113    | 102    | 215     |       | 22    | 237   |
| (%)          | (52.6) | (47.4) | (100.0) |       |       |       |
| 14～          | 40     | 33     | 73      |       | 6     | 79    |
| (%)          | (54.8) | (45.2) | (100.0) |       |       |       |
| 合計           | 3,173  | 2,314  | 5,487   | 21    | 439   | 5,947 |
| (%)          | (57.8) | (42.2) | (100.0) |       |       |       |
| 記載なし         | 500    | 938    | 1,438   | 5     | 1,551 | 2,994 |
| (%)          | (34.8) | (65.2) | (100.0) |       |       |       |
| 総計           | 3,673  | 3,252  | 6,925   | 26    | 1,990 | 8,941 |
| (%)          | (53.0) | (47.0) | (100.0) |       |       |       |
| 平均           | 10.83  | 10.79  | 10.81   | 10.35 | 10.85 | 10.81 |
| 標準偏差         | 1.36   | 1.50   | 1.42    | 1.27  | 1.40  | 1.42  |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査