

表 693 糖尿病の既往 Non-HDL-C濃度(mg/dL)別（腹膜透析患者）

Non-HDL-C濃度	既往なし	既往あり	合計	不明	記載なし	総計
50未満	8	9	17		1	18
(%)	(47.1)	(52.9)	(100.0)			
50～	37	70	107		7	114
(%)	(34.6)	(65.4)	(100.0)			
70～	166	160	326	1	24	351
(%)	(50.9)	(49.1)	(100.0)			
90～	316	300	616	2	42	660
(%)	(51.3)	(48.7)	(100.0)			
110～	436	301	737	1	59	797
(%)	(59.2)	(40.8)	(100.0)			
130～	393	215	608	1	53	662
(%)	(64.6)	(35.4)	(100.0)			
150～	245	157	402	1	33	436
(%)	(60.9)	(39.1)	(100.0)			
170～	153	90	243		20	263
(%)	(63.0)	(37.0)	(100.0)			
190～	64	49	113		7	120
(%)	(56.6)	(43.4)	(100.0)			
210～	49	34	83		7	90
(%)	(59.0)	(41.0)	(100.0)			
合計	1,867	1,385	3,252	6	253	3,511
(%)	(57.4)	(42.6)	(100.0)			
記載なし	1,806	1,867	3,673	20	1,737	5,430
(%)	(49.2)	(50.8)	(100.0)			
総計	3,673	3,252	6,925	26	1,990	8,941
(%)	(53.0)	(47.0)	(100.0)			
平均	131.21	124.57	128.38	115.50	129.91	128.47
標準偏差	36.99	39.10	38.04	27.33	37.94	38.01

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査