

表 715 心筋梗塞の既往 身長(cm)別（透析患者全体）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	277	33	310	2	42	354
(%)	(89.4)	(10.6)	(100.0)			
130～	542	51	593	2	101	696
(%)	(91.4)	(8.6)	(100.0)			
135～	2,171	192	2,363	19	455	2,837
(%)	(91.9)	(8.1)	(100.0)			
140～	7,877	673	8,550	48	1,699	10,297
(%)	(92.1)	(7.9)	(100.0)			
145～	17,948	1,663	19,611	65	3,997	23,673
(%)	(91.5)	(8.5)	(100.0)			
150～	28,839	2,724	31,563	107	6,565	38,235
(%)	(91.4)	(8.6)	(100.0)			
155～	33,297	3,669	36,966	129	7,377	44,472
(%)	(90.1)	(9.9)	(100.0)			
160～	39,333	5,185	44,518	138	8,994	53,650
(%)	(88.4)	(11.6)	(100.0)			
165～	35,257	4,741	39,998	104	7,886	47,988
(%)	(88.1)	(11.9)	(100.0)			
170～	22,965	2,749	25,714	87	5,273	31,074
(%)	(89.3)	(10.7)	(100.0)			
175～	8,527	859	9,386	24	1,909	11,319
(%)	(90.8)	(9.2)	(100.0)			
180～	2,290	180	2,470	9	500	2,979
(%)	(92.7)	(7.3)	(100.0)			
185～	366	20	386		79	465
(%)	(94.8)	(5.2)	(100.0)			
190～	57	6	63		14	77
(%)	(90.5)	(9.5)	(100.0)			
合計	199,746	22,745	222,491	734	44,891	268,116
(%)	(89.8)	(10.2)	(100.0)			
記載なし	12,342	1,451	13,793	183	28,016	41,992
(%)	(89.5)	(10.5)	(100.0)			
総計	212,088	24,196	236,284	917	72,907	310,108
(%)	(89.8)	(10.2)	(100.0)			
平均	159.96	160.81	160.05	158.58	160.04	160.05
標準偏差	9.71	9.12	9.66	10.62	9.61	9.65

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査