

表 724 心筋梗塞の既往 透析前ヘモグロビン濃度(g/dL)別（透析患者全体）

透析前ヘモグロビン濃度	既往なし	既往あり	合計	不明	記載なし	総計
5未満	242	28	270		33	303
(%)	(89.6)	(10.4)	(100.0)			
5～	335	26	361	2	66	429
(%)	(92.8)	(7.2)	(100.0)			
6～	843	96	939	15	206	1,160
(%)	(89.8)	(10.2)	(100.0)			
7～	3,287	384	3,671	19	745	4,435
(%)	(89.5)	(10.5)	(100.0)			
8～	10,824	1,341	12,165	59	2,330	14,554
(%)	(89.0)	(11.0)	(100.0)			
9～	34,257	3,932	38,189	130	7,670	45,989
(%)	(89.7)	(10.3)	(100.0)			
10～	70,134	7,644	77,778	275	15,859	93,912
(%)	(90.2)	(9.8)	(100.0)			
11～	58,452	6,378	64,830	176	13,167	78,173
(%)	(90.2)	(9.8)	(100.0)			
12～	21,510	2,468	23,978	80	4,796	28,854
(%)	(89.7)	(10.3)	(100.0)			
13～	5,410	676	6,086	10	1,212	7,308
(%)	(88.9)	(11.1)	(100.0)			
14～	1,984	293	2,277	4	392	2,673
(%)	(87.1)	(12.9)	(100.0)			
合計	207,278	23,266	230,544	770	46,476	277,790
(%)	(89.9)	(10.1)	(100.0)			
記載なし	4,810	930	5,740	147	26,431	32,318
(%)	(83.8)	(16.2)	(100.0)			
総計	212,088	24,196	236,284	917	72,907	310,108
(%)	(89.8)	(10.2)	(100.0)			
平均	10.71	10.72	10.71	10.46	10.71	10.71
標準偏差	1.28	1.32	1.29	1.38	1.26	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査