

表 726 心筋梗塞の既往 HDL-C濃度(mg/dL)別 (透析患者全体)

HDL-C濃度	既往なし	既往あり	合計	不明	記載なし	総計
20未満	570	92	662	2	132	796
(%)	(86.1)	(13.9)	(100.0)			
20～	11,193	1,905	13,098	41	2,484	15,623
(%)	(85.5)	(14.5)	(100.0)			
30～	33,450	4,841	38,291	139	7,417	45,847
(%)	(87.4)	(12.6)	(100.0)			
40～	38,386	4,306	42,692	136	8,524	51,352
(%)	(89.9)	(10.1)	(100.0)			
50～	28,350	2,696	31,046	102	6,413	37,561
(%)	(91.3)	(8.7)	(100.0)			
60～	17,145	1,273	18,418	53	3,765	22,236
(%)	(93.1)	(6.9)	(100.0)			
70～	8,480	558	9,038	30	1,924	10,992
(%)	(93.8)	(6.2)	(100.0)			
80～	4,031	256	4,287	14	928	5,229
(%)	(94.0)	(6.0)	(100.0)			
90～	1,772	94	1,866	5	460	2,331
(%)	(95.0)	(5.0)	(100.0)			
100～	1,471	109	1,580	3	418	2,001
(%)	(93.1)	(6.9)	(100.0)			
合計	144,848	16,130	160,978	525	32,465	193,968
(%)	(90.0)	(10.0)	(100.0)			
記載なし	67,240	8,066	75,306	392	40,442	116,140
(%)	(89.3)	(10.7)	(100.0)			
総計	212,088	24,196	236,284	917	72,907	310,108
(%)	(89.8)	(10.2)	(100.0)			
平均	49.05	44.65	48.61	47.68	49.47	48.75
標準偏差	16.66	15.35	16.59	15.58	17.27	16.71

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査